arXiv:1302.6627v2 [math.RT] 6 Mar 2013

REGULAR FUNCTIONS OF NILPOTENT ORBITS AND NORMALITY OF THEIR CLOSURES

KAYUE (DANIEL) WONG

Abstract. For any nilpotent orbit $\mathcal{O}$ in a complex semisimple Lie group G, it is known from algebraic geometry that the normality of its Zariski closure $\overline{\mathcal{O}}$ is related to the ring of regular functions of $\mathcal{O}$. On the other hand, if G is classical, Kraft and Procesi gave a combinatorial criterion on the normality of $\overline{\mathcal{O}}$. In this paper, we will give a algorithm computing the multiplicities of the fundamental representations in the ring of regular functions of $\mathcal{O}$, and will relate it to an equivalent criterion on the normality of $\overline{\mathcal{O}}$ in terms of multiplicities of fundamental representations.

1. Preliminaries

1.1. **Classical Nilpotent Orbits.** It is well-known that all nilpotent orbits in a complex classical Lie group can be expressed as partitions, where the partition corresponds to the size of the Jordan blocks. In particular, here is the classification of nilpotent orbits in type B, C and D, given in [6] for instance:

Theorem 1.1. *For type B_n, all nilpotent orbits in $SO(2n+1,\mathbb{C})$ are identified with the partitions of $2n+1$ in which even parts occur with even multiplicity.*

For type C_n, nilpotent orbits in $Sp(2n,\mathbb{C})$ are identified with the paritions of $2n$ in which odd parts occur with even multiplicity.

For type D_n, nilpotent orbits in $SO(2n,\mathbb{C})$ are identified with the partitions of $2n$ in which even parts occurs with even multiplicity, except for the 'very even' partitions - partitions with only even parts, each having even multiplicity. Each very even partitions corresponds to 2 orbits.

If we only consider $O(2n,\mathbb{C})$-conjugates of nilpotent elements in $\mathfrak{o}(2n,\mathbb{C}) = \mathfrak{so}(2n,\mathbb{C})$, the two nilpotent orbits corresponding to a very even partition will be merged into one orbit. In this case, there is at most one orbit for every partition of $2n$.

From now on, we only consider $O(2n,\mathbb{C})$-conjugates of nilpotent elements in $\mathfrak{so}(2n,\mathbb{C})$. Therefore, there is only one orbit for the very even partitions of $2n$.

1.2. **Another Characterization of Classical Nilpotent Orbits.** In the last Section, all nilpotent orbits of classical type are characterized by partitions. And the partitions are often expressed as Young diagrams whose row sizes are determined by the corresponding partitions. In fact, in studying nilpotent orbits, it is sometimes more convenient to look at the column sizes of a Young diagram. The column sizes of the Young diagram corresponding to a partition is given by the **dual partition** of the original partition, which is defined by the following:

Definition 1.2. *Let* $(r_1, r_2, \ldots, r_i)$ *be a partition of* n*, with* $r_1 \geq r_2 \geq \cdots \geq r_i > 0$*, then its* ***dual partition*** *is given by* $(c_k, c_{k-1}, \ldots, c_1)$*, where* $c_{k+1-j} = \#\{i | r_i \geq j\}$*.*

Example 1.3. *Let* $\mathcal{O} = [4, 2]$ *in* $Sp(6, \mathbb{C})$*. Then the Young diagram corresponding to* $\mathcal{O}$ *is given by*



And the dual partition of $\mathcal{O} = (2, 2, 1, 1)$*.*

From now on, we will determine a nilpotent orbit by its dual partition, or equivalently the column sizes of its corresponding Young diagram. Here is a restatement of Theorem 1.1 in terms of column sizes.

Corollary 1.4. *Any nilpotent orbit in* $Sp(2m, \mathbb{C})$ *can be paramatrized by a partition of* $2m$ *with column sizes* $(c_{2k}, c_{2k-1}, \ldots, c_0)$*, where* $c_{2k} \geq c_{2k-1} \geq \cdots \geq c_0 \geq 0$ *(by insisting* c_{2k} *is the longest column, we put* $c_0 = 0$ *if necessary), such that* $c_{2i} + c_{2i-1}$ *is even for all* $i \geq 0$*. (Note:* $c_{-1} = 0$*)*
Any nilpotent orbit in $O(n, \mathbb{C})$ *can be paramatrized by a partition of* n *with column sizes* $(b_{2k+1}, b_{2k}, \ldots, b_0)$*, where* $b_{2k+1} \geq b_{2k} \geq \cdots \geq b_0 \geq 0$ *(putting* $b_0 = 0$ *if necessary), such that* $b_{2i} + b_{2i-1}$ *is even for all* $i \geq 0$*. (Note:* $b_{-1} = 0$*)*

Example 1.5. $(7, 6, 4, 4)$ *is a nilpotent orbit in* $O(21, \mathbb{C})$ *but it does not define a nilpotent orbit in* Sp*.*

1.3. **Kraft-Procesi Criterion on Normality.** Given the combinatorial description of the nilpotent orbit $\mathcal{O}$ in the classical case, we can now state the criterion on the normality of the Zariski closure of $\mathcal{O}$, given by Kraft and Procesi:

Theorem 1.6 (Kraft-Procesi)**.** *Let* $\mathcal{O} = (c_{2k}, c_{2k-1}, \ldots, c_0)$ *be a nilpotent orbit in* $Sp(2m, \mathbb{C})$*. If there is a chain of column lengths of the form*

$$c_{2i} \neq c_{2i-1} = c_{2i-2} = \cdots = c_{2j-1} = c_{2j-2} \neq c_{2j-3}$$

then $\overline{\mathcal{O}}$ *is not normal along* $(c_{2k}, \ldots, c_{2i}, c_{2i-1}+2, c_{2i-2}, \ldots, c_{2j-1}, c_{2j-2}-2, c_{2j-3}, \ldots, c_0)$*.*
Similarly, the closure of a nilpotent orbit $\mathcal{P} = (b_{2k+1}, \ldots, b_0)$ *in* $O(n, \mathbb{C})$ *is not normal if there is a chain of column lengths of the form*

$$b_{2i} \neq b_{2i-1} = b_{2i-2} = \cdots = b_{2j-1} = b_{2j-2} \neq b_{2j-3}$$

(Note: $b_{2k+2} = 0$*)*

Example 1.7. *In* $Sp(2m, \mathbb{C})$*, the orbit closures* $\overline{(8,6,6,6)}$*,* $\overline{(6,6,6,6)}$ *are normal, while* $\overline{(8,6,6,4)}$ *is not normal. In* $O(n, \mathbb{C})$*,* $\overline{(8,6,6,6)}$*,* $\overline{(8,6,6,4)}$ *are normal,* $\overline{(6,6,6,6)}$ *is not normal.*

2. Statement of the Main Results

Apart from the combinatorial criterion of normality given in the last Section, there is a more general algebro-geometric criterion of normality:

Lemma 2.1. *Let* $\mathcal{O}$ *be a* G*-orbit with Zariski closure* $\overline{\mathcal{O}}$*. If* $\overline{\mathcal{O}} \setminus \mathcal{O}$ *has codimension greater than or equal to 2, then the ring of regular functions of* $\mathcal{O}$*,* $R[\mathcal{O}]$ *is the integral closure of* $R[\overline{\mathcal{O}}]$ *in its field of fractions.*

Note that all (real or complex) nilpotent orbits are symplectic manifolds with the **Kostant-Sekiguchi symplectic form**, therefore they are all of even (real or complex) dimensions. In particular, the nilpotent orbit closure $\overline{\mathcal{O}}$ satisfies the hypothesis of the above Theorem. So we have the following:

Corollary 2.2. $R[\mathcal{O}] \cong R[\overline{\mathcal{O}}]$ *if and only if* $\overline{\mathcal{O}}$ *is normal.*

Also, it is well-known that if X is an algebraic variety endowed with an action of a reductive group G (so $\mathcal{O}$ and $\overline{\mathcal{O}}$ are examples of X), then $R[X]$ can be decompsed as a direct sum of irreducible, finite-dimensional G representations, i.e. $R[X] = \bigoplus_\lambda m_\lambda V_\lambda$ for λ being the parameters of irreducible, finite-dimensional G representations. The first Theorem of the paper is to compute the multiplicities m_λ for fundamental representations V_λ in $R[\mathcal{O}]$:

Theorem 2.3. *Suppose* $\mathcal{O} = (c_{2k}, c_{2k-1}, \dots, c_0)$ *is a nilpotent orbit in* $Sp(2m, \mathbb{C})$*. First remove all column pairs of same size, leaving the orbit* $(d_{2l}, d_{2l-1}, \dots, d_0)$*. For each removed column pair* $c_i = c_{i-1} = y$*, let* $Y = \{c_i | c_i = c_{i-1} \text{ are removed from } \mathcal{O}\}$ *with multiplicities. Also, let* $Z = \{z_j = \frac{d_{2j}+d_{2j-1}}{2} | j = 0, \dots, l\}$*. Then rearrange elements in* $Y \cup Z$ *in non-decreasing order to get* $W := Y \cup Z = \{w_i | i = 0, 1, \dots, k\}$*, with* $w_i \leq w_j$ *if* $i < j$*.*

Now define a sequence of sequences $\beta_i = (\beta_{i0}, \beta_{i1}, \dots)$ *recursively by:*

- *Begin with the sequence* $\beta_0 = (\beta_{00}, \beta_{01}, \beta_{02}, \dots) = (\binom{k-1}{0}, \binom{k}{1}, \binom{k+1}{2}, \dots)$*.*
- *Define the* i*-th sequence* α_i *recursively by* $\beta_{i+1} = \beta_i - (\overbrace{0, \dots, 0}^{w_i+1}, \beta_{i0}, \beta_{i1}, \beta_{i2}, \dots)$*.*

Then

$$[R[\mathcal{O}] : \mu_{2i}] = \beta_{(k+1)i} \text{ for } i \leq \frac{m}{2}$$

$$[R[\mathcal{O}] : \mu_{2i+1}] = 0 \text{ for all } i$$

where $\mu_i = \wedge^i\mathbb{C}^{2n}/\wedge^{i-2}\mathbb{C}^{2n}$ *are the fundamental representations of* $Sp(2n,\mathbb{C})$ *(if* $i-2<0$*, take* $\wedge^{i-2}\mathbb{C}^{2n} = triv$*).*

Suppose $\mathcal{P} = (c_{2k+1}, c_{2k}, c_{2k-1}, \dots, c_0)$ *is a nilpotent orbit in* $O(n,\mathbb{C})$*. First remove all column pairs of same size, leaving the orbit* $(d_{2l+1}, d_{2l}, d_{2l-1}, \dots, d_0)$*. For each removed column pair* $c_i = c_{i-1} = y$*, let* $Y = \{y | c_i = c_{i-1} \text{ are removed from } \mathcal{P}\}$ *with multiplicities. Also, let* $Z = \{z_j = \frac{d_{2j}+d_{2j-1}}{2} | j = 0,\dots,l\}$ *(note that* d_{2l+1} *is not used in the algorithm). Then rearrange elements in* $Y \cup Z$ *in non-decreasing order to get* $Y \cup Z = \{w_i | i = 0,1,\dots,k\}$*, with* $w_i \leq w_j$ *if* $i<j$*.*
Now define a sequence of sequences $\beta_i = (\beta_{i0}, \beta_{i1}, \dots)$ *recursively by:*

- *Begin with the sequence* $\beta_0 = (\beta_{00}, \beta_{01}, \beta_{02}, \dots) = (\binom{k}{0}, \binom{k+1}{1}, \binom{k+2}{2}, \dots)$*.*
- *Define the i-th sequence* α_i *recursively by* $\beta_{i+1} = \beta_i - (\overbrace{0,\dots,0}^{w_i+1}, \beta_{i0}, \beta_{i1}, \beta_{i2}, \dots)$*.*

Then for the fundamental representations $\mu'_j := \wedge^j\mathbb{C}^n$ *in* $O(n,\mathbb{C})$*,*

$$[R[\mathcal{P}] : \mu'_{2i}] = \beta_{(k+1)i} \text{ for } 2i \leq n$$

$$[R[\mathcal{P}] : \mu'_{2i+1}] = 0 \text{ for all } i$$

Example 2.4. *Let* $\mathcal{O} = (8,6,6,4,4,2,2)$ *in* $Sp(32,\mathbb{C})$*. Then the* w_i*'s are* $\{2,4,4,6\}$*. We therefore have the multiplicities as follows:*

i	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
$[R[\mathcal{O}] : \mu_i]$	1	0	3	0	6	0	9	0	12	0	13	0	12	0	8	0	3

Let $\mathcal{P} = (7,5,3,3,1)$ *in* $O(19,\mathbb{C})$*. Then the* w_i*'s are* $\{6,6\}$*. We therefore have the multiplicities of* μ'_{2i} *as follows:*

i	0	2	4	6	8	10	12	14	16	18
$[R[\mathcal{P}] : \mu'_i]$	1	2	3	4	3	2	1	0	0	0

The second Theorem of this paper is to relate the algebro-geometric criterion of normality in Corollary 2.2 with the combinatorial criterion given by Kraft-Procesi in Theorem 1.6. In particular, we have the following Theorem:

Corollary 2.5. *Let* $\mathcal{O}$ *be a nilpotent orbit in* $Sp(2m,\mathbb{C})$*. Then* $\overline{\mathcal{O}}$ *is not normal iff*

$$[R[\overline{\mathcal{O}}] : \mu_i] < [R[\mathcal{O}] : \mu_i]$$

for some $i > 0$*.*
Let $\mathcal{P} = (b_{2k+1}, \dots, b_0)$ *be a nilpotent orbit in* $O(n,\mathbb{C})$ *such that* $b_{2k+1} \neq b_{2k}$*. Then*

$\overline{\mathcal{P}}$ is not normal iff

$$[R[\overline{\mathcal{P}}] : \mu_i'] < [R[\mathcal{P}] : \mu_i']$$

for some $i > 0$.

Remark 2.6. *In fact, the Corollary holds for all orbits in $O(n, \mathbb{C})$ without the condition $b_{2k+1} \neq b_{2k}$. However, the proof will be a lot more involved in this special case, which we do not include here.*

3. Proof of Theorem 2.3

3.1. **The Barbasch Model on $\mathcal{O}$.**

In [4, Section 2], Barbasch constructed a $(\mathfrak{g}_{\mathbb{C}}, K_{\mathbb{C}})$-module of any nilpotent orbit of classical type, denoted as $X_{\mathcal{O}}$. The definition of $X_{\mathcal{O}}$ is given below:

Definition 3.1. *Let $\mathcal{O} = (c_r, \dots, c_0)$ be any nilpotent orbit in G, where $G = Sp(2m, \mathbb{C})$ or $O(n, \mathbb{C})$. Define Y and $\mathcal{Q}$ inductively as follows:*
Let $Y_0 = \phi$ and $\mathcal{Q}_0 = \mathcal{O}$. Suppose i is the smallest integer such that $c_i = c_{i+1}$, then $\mathcal{Q}_1 := (c_r, \dots, c_{i+2}, c_{i-1}, \dots, c_0)$ and $Y_1 := Y_0 \cup \{c_i\}$.

Continue the above process until we get $\mathcal{Q} = (d_l, \dots, d_0)$, with $d_i \neq d_{i+1}$ for all i. Then Y are the lengths of the removed columns with multiplicities. Let

$$X_{\mathcal{O}} := Ind_{G(\mathcal{Q},\mathbb{C}) \times GL(Y)}^{G(2m,\mathbb{C})} (X_{\mathcal{Q}} \otimes triv \otimes \cdots \otimes triv) \tag{1}$$

*where $G(\mathcal{Q}, \mathbb{C}) = G(\sum d_i, \mathbb{C})$, $GL(Y) = \Pi_{c_i \in Y} GL(c_i, \mathbb{C})$ and $X_{\mathcal{Q}}$ is the **spherical unipotent representation** attached to the nilpotent orbit $\mathcal{Q}$.*

Theorem 3.2 (Barbasch)**.** *As $K_{\mathbb{C}} \cong G$-modules,*

$$X_{\mathcal{O}} \cong R[\mathcal{O}]$$

The theory of unipotent representation is well-studied by Barbasch and Vogan, for example [2], [3]. In particular, the character theory of such representations are known. This gives us in theory a way to compute the multiplicities of any finite-dimensional irreducible representations of $X_{\mathcal{Q}}$.

3.2. **Spherical Unipotent Representations.**

In this Subsection, we explore the theory of spherical unipotent representations mentioned in the last Subsection.

Proposition 3.3. *Let $\mathcal{Q} = (c_{2k}, \dots, c_0)$ be a nilpotent orbit in $Sp(2n, \mathbb{C})$, with $c_i \neq c_{i-1}$ for all i. Then the corresponding infinitesimal character attached to $\mathcal{Q}$ is $(\chi; \chi) = (\chi_0, \chi_1, \dots, \chi_k; \chi_0, \chi_1, \dots, \chi_k)$, where χ_i is defined by the following:*

- *For i between 1 and k, $\chi_i = (\frac{c_{2i}}{2}, \frac{c_{2i}-2}{2}, \dots, \frac{-c_{2i-1}+2}{2})$*
- *$\chi_0 = (\frac{c_0}{2}, \dots, 1)$*

Let $\mathcal{P} = (b_{2k+1}, b_{2k}, b_{2k-1}, \dots b_0)$ be any nilpotent orbit in $O(n, \mathbb{C})$. The infinitesimal character of $X_{\mathcal{P}}$ is $(\chi, \chi_{k+1}; \chi, \chi_{k+1})$, where χ is defined in the same way for $\mathcal{P}' = (b_{2k}, b_{2k-1}, \dots, b_0)$, and $\chi_{k+1} = (\frac{b_{2k+1}-2}{2}, \dots, \frac{1}{2})$ if b_{2k+1} is odd, $(\frac{b_{2k+1}-2}{2}, \dots, 0)$ if b_{2k+1} is even.

Proof. This is given in [3]. □

Therefore, the character formula of $X_{\mathcal{Q}}$ in the last Subsection consists of integral and half-integral entries. We first study the character theory in these two separate cases.

3.2.1. *Integral Character Theory.* Suppose the infinitesimal character is integral, then it is necessary that all the columns sizes of $\mathcal{Q}$ are even. We need a few notations before stating the character formula of $X_{\mathcal{Q}}$.

Definition 3.4. *Let $\chi \in \mathfrak{h}^*$ be an integral character. Correspond $\mathfrak{h}^*$ with the Langlands L-group ${}^L\mathfrak{h}$. Next, find the Levi subalgebra ${}^L\mathfrak{m} \subset^L \mathfrak{h}$ such that $\langle \lambda, m \rangle = 0$ for all $m \in^L \mathfrak{m}$. Then the* ***left cell*** *corresponding to χ is given by $V(\chi) = J_{W({}^L\mathfrak{m})}^W(sgn) \otimes sgn$, where J is the truncated induction defined by Lusztig.*

Theorem 3.5. *Let $\mathcal{Q} = (d_k, \dots, d_1)$ be a nilpotent orbit in $\mathfrak{g}$ with $d_i \neq d_{i-1}$ for all i, and let χ, $V(\chi)$ as in the previous Sections. Then there is a one-one correspondence between the irreducible Weyl group representations in $V(\chi)$ and $(\mathbb{Z}/2\mathbb{Z})^s$, where $s = \lfloor \frac{k}{2} \rfloor$ if $\mathfrak{g} = \mathfrak{sp}$, and $s = \lceil \frac{k}{2} \rceil - 1$ if $\mathfrak{g} = \mathfrak{o}$. Suppose the correspondence is given by*

$$x \in (\mathbb{Z}/2\mathbb{Z})^s \longleftrightarrow \sigma_x \in V(\chi)$$

then the character formula of $X_{\mathcal{Q}}$ is

$$X_{\mathcal{Q}} = \frac{1}{2^s} \sum_x R_{\sigma_x}(\chi)$$

with

$$R_\sigma(\chi) := \sum_{w \in W} tr(\sigma(w)) X \begin{pmatrix} \chi \\ w\chi \end{pmatrix}$$

where $X \begin{pmatrix} \lambda \\ \mu \end{pmatrix}$ is the K-finite part of the principal series representation $Ind_B^G(\mathbb{C}_{(\lambda,\mu)})$ ([2, Definition 1.7(d)]*)*

By using the techniques in [2, Chapter 6], the above character formula can be substantially simplified. Here is an example of how the formula of $X_{\mathcal{Q}}$ is like:

Example 3.6. *Suppose $\mathcal{Q} = (8,6,4,2)$ in $Sp(20,\mathbb{C})$. Then $\chi = (4,3,2,1,0,1,2;2,1,0)$. The character formula of $X_{\mathcal{Q}}$ is*

$$\frac{1}{4}\sum_{W(C_4\times D_3\times C_2\times D_1)}(-1)^{l(w)}X\left(\begin{array}{ccc} & 4321,210 & 21,0 \\ w(& 4321,210 & 21,0) \end{array}\right)$$

$$+\frac{1}{4}\sum_{W(D_5\times C_2\times C_2\times D_1)}(-1)^{l(w)}X\left(\begin{array}{ccc} & 43210,21 & 21,0 \\ w(& 43210,21 & 21,0) \end{array}\right)$$

$$+\frac{1}{4}\sum_{W(C_4\times D_3\times D_3\times C_0)}(-1)^{l(w)}X\left(\begin{array}{ccc} & 4321,210 & 210 \\ w(& 4321,210 & 210) \end{array}\right)$$

$$+\frac{1}{4}\sum_{W(D_5\times C_2\times D_3\times C_0)}(-1)^{l(w)}X\left(\begin{array}{ccc} & 43210,21 & 210 \\ w(& 43210,21 & 210) \end{array}\right)$$

Let $\mathcal{P} = (6,4,2)$ in $O(12,\mathbb{C})$, the character formula of $X_{\mathcal{P}}$ is

$$\frac{1}{2}\sum_{W(C_2\times D_1)}(-1)^{l(w)}X\left(\begin{array}{ccc} & 210+ & 21,0 \\ w(& 210+ & 21,0) \end{array}\right)+\frac{1}{2}\sum_{W(D_3\times C_0)}(-1)^{l(w)}X\left(\begin{array}{ccc} & 210+ & 210 \\ w(& 210+ & 210) \end{array}\right)$$

3.2.2. *Half-Integral Character Theory.* After dealing with the case of purely integral infinitesimal characters, we move to the case of purely half-integral infinitesimal characters. This is an important ingredient for such theory:

Theorem 3.7 (Kazhdan-Lusztig Conjecture). *For each $\lambda \in^L \mathfrak{h}$, consider the root system $\Delta(\lambda) := \{\alpha | \langle\check{\alpha}, \lambda\rangle \in \mathbb{Z}\}$. Then the character theory of unipotent representations in $\mathfrak{g}$ can be derived from that of $\Delta(\lambda)$.*

The conjecture was proved separately by Kashiwara-Vergne and Beilinson- Bernstein.

By the Kazhdan-Lusztig conjecture, we study the character theory of unipotent representations with half-integral infinitesimal characters in $W(D_m) \leq W(C_m)$ (note that the Kazhdan-Lusztig conjecture is vacuous for half-integral infinitesimal characters in the $W(D_n)$ and $W(B_n)$ cases).

In this case, the character formula is different from the integral case. It is better expressed by examples:

Example 3.8. *Let $\mathcal{Q} = (9,7)$ in $Sp(16,\mathbb{C})$. Then $\chi = (\frac{9}{2},\frac{7}{2},\frac{5}{2},\frac{3}{2},\frac{1}{2},\frac{-1}{2},\frac{-3}{2},\frac{-5}{2})$, and the character formula of $\mathcal{Q}$ is*

$$\sum_{w\in W(D_5\times D_3)}(-1)^{l(w)}X\left(\begin{array}{ccc} & \frac{9}{2}\frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2} \\ w(& \frac{9}{2}\frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2}) \end{array}\right)-\sum_{w\in W(D_5\times D_3)}(-1)^{l(w)}X\left(\begin{array}{ccc} & \frac{9}{2}\frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2} \\ w(& \frac{9}{2}\frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{-1}{2} & \frac{5}{2}\frac{3}{2}\frac{-1}{2}) \end{array}\right)$$

Let $\mathcal{P} = (9,7,5,3)$ in $Sp(24,\mathbb{C})$, the character formula of $X_{\mathcal{P}}$ is:

$$\sum_{w\in W(D_5\times D_3\times D_3\times D_1)} (-1)^{l(w)} X\begin{pmatrix} & \frac{9}{2}\frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{1}{2} \\ w(& \frac{9}{2}\frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{1}{2}) \end{pmatrix}$$

$$- \sum_{w\in W(D_5\times D_3\times D_3\times D_1)} (-1)^{l(w)} X\begin{pmatrix} & \frac{9}{2}\frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{1}{2} \\ w(& \frac{9}{2}\frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{-1}{2} & \frac{5}{2}\frac{3}{2}\frac{-1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{1}{2}) \end{pmatrix}$$

$$- \sum_{w\in W(D_5\times D_3\times D_3\times D_1)} (-1)^{l(w)} X\begin{pmatrix} & \frac{9}{2}\frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{1}{2} \\ w(& \frac{9}{2}\frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{-1}{2} & \frac{-1}{2}) \end{pmatrix}$$

$$+ \sum_{w\in W(D_5\times D_3\times D_3\times D_1)} (-1)^{l(w)} X\begin{pmatrix} & \frac{9}{2}\frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{1}{2} \\ w(& \frac{9}{2}\frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{-1}{2} & \frac{5}{2}\frac{3}{2}\frac{-1}{2} & \frac{5}{2}\frac{3}{2}\frac{-1}{2} & \frac{-1}{2}) \end{pmatrix}$$

3.2.3. *General Case.* In the previous subsections, we have seen the character theory of unipotent represenations with purely integral or purely half-integral infinitesimal characters. In fact, by the Kazhdan-Lusztig Conjecture (Theorem 3.7), it is enough to derive the theory of unipotent represenations out of these two cases. One just need to separate the integral and half-integral coordinates, apply the algorithms in the previous two sections, combine them together to get the result.

Example 3.9. *Let $\mathcal{P} = (9,7,5,4,2)$ in $O(27,\mathbb{C})$. Then from above, we know the character formula of the orbit $(9,7,5)$ in $O(21,\mathbb{C})$ is*

$$\sum_{W(D_4\times D_2)} (-1)^{l(w)} X\begin{pmatrix} & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2}+ & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{3}{2}\frac{1}{2} \\ w(& \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2}+ & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{3}{2}\frac{1}{2}) \end{pmatrix}$$

$$- \sum_{W(D_4\times D_2)} (-1)^{l(w)} X\begin{pmatrix} & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2}+ & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{3}{2}\frac{1}{2} \\ w(& \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2}+ & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{-1}{2} & \frac{3}{2}\frac{-1}{2}) \end{pmatrix}$$

and the character formula for the orbit $(4,2)$ in $Sp(6,\mathbb{C})$ is

$$\frac{1}{2}\Big(\sum_{W(C_2\times D_1)} (-1)^{l(w)} X\begin{pmatrix} & 21 & 0 \\ w(& 21 & 0) \end{pmatrix} + \sum_{W(D_3\times C_0)} (-1)^{l(w)} X\begin{pmatrix} & 210 \\ w(& 210) \end{pmatrix}\Big)$$

Then the character formula for the $X_{\mathcal{P}}$ is just the 'cocatenation' of the two character formulas above:

$$\frac{1}{2} \sum_{W(D_4\times D_2\times C_2\times D_1)} (-1)^{l(w)} X\begin{pmatrix} & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2}+ & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{3}{2}\frac{1}{2} & 21 & 0 \\ w(& \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2}+ & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{3}{2}\frac{1}{2} & 21 & 0) \end{pmatrix}$$

$$-\frac{1}{2}\sum_{W(D_4\times D_2\times C_2\times D_1)}(-1)^{l(w)}X\left(\begin{array}{cccccc} & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2}+ & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{3}{2}\frac{1}{2} & 21 & 0 \\ w(& \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2}+ & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{-1}{2} & \frac{3}{2}\frac{-1}{2} & 21 & 0) \end{array}\right)$$

$$+\frac{1}{2}\sum_{W(D_4\times D_2\times D_3\times C_0)}(-1)^{l(w)}X\left(\begin{array}{ccccc} & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2}+ & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{3}{2}\frac{1}{2} & 210 \\ w(& \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2}+ & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{3}{2}\frac{1}{2} & 210) \end{array}\right)$$

$$-\frac{1}{2}\sum_{W(D_4\times D_2\times D_3\times C_0)}(-1)^{l(w)}X\left(\begin{array}{ccccc} & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2}+ & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2} & \frac{3}{2}\frac{1}{2} & 210 \\ w(& \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{1}{2}+ & \frac{7}{2}\frac{5}{2}\frac{3}{2}\frac{-1}{2} & \frac{3}{2}\frac{-1}{2} & 210) \end{array}\right)$$

In fact, the above cocantenation technique works in general:

Proposition 3.10. *If p,q are both even, define $\overbrace{p,q}^{+}$ to be*

$$\frac{1}{2}\sum_{W(C_{p/2}\times D_{q/2+1})}(-1)^{l(w)}X\left(\begin{array}{ccc} & \frac{p}{2}\ldots 1 & \frac{q}{2}\ldots 0 \\ w(& \frac{p}{2}\ldots 1 & \frac{q}{2}\ldots 0) \end{array}\right)$$

$$+\frac{1}{2}\sum_{W(D_{p/2+1}\times C_{q/2})}(-1)^{l(w)}X\left(\begin{array}{ccc} & \frac{p}{2}\ldots 0 & \frac{q}{2}\ldots 1 \\ w(& \frac{p}{2}\ldots 0 & \frac{q}{2}\ldots 1) \end{array}\right))$$

If p,q are both odd, define $\overbrace{p,q}^{+}$ to be

$$\sum_{W(D_{p/2}\times D_{q/2-1})}(-1)^{l(w)}X\left(\begin{array}{ccc} & \frac{p}{2}\cdots\frac{1}{2} & \frac{q-2}{2}\cdots\frac{1}{2} \\ w(& \frac{p}{2}\cdots\frac{1}{2} & \frac{q-2}{2}\cdots\frac{1}{2}) \end{array}\right)$$

$$-\sum_{W(D_{p/2}\times D_{q/2-1})}(-1)^{l(w)}X\left(\begin{array}{ccc} & \frac{p}{2}\cdots\frac{3}{2}\frac{1}{2} & \frac{q-2}{2}\cdots\frac{3}{2}\frac{1}{2} \\ w(& \frac{p}{2}\cdots\frac{3}{2}\frac{-1}{2} & \frac{q-2}{2}\cdots\frac{3}{2}\frac{-1}{2}) \end{array}\right)$$

Let $\mathcal{O} = (d_{2l},\ldots,c_0)$ be an orbit in $Sp(2m,\mathbb{C})$ so that no two columns are of the same size. Then the character formula of $X_{\mathcal{O}}$ is of the form

$$X_{\mathcal{O}} \cong (\overbrace{d_{2l},d_{2l-1}}^{+};\overbrace{d_{2l-2},d_{2l-3}}^{+};\ldots;\overbrace{d_2,d_1}^{+};d_0)$$

i.e. the cocatenation of the character formulas $\overbrace{d_{2l},d_{2l-1}}^{+}$, $\overbrace{d_{2l-1},d_{2l-2}}^{+}$, $\ldots$, d_0, and d_0 has the character formula $T := \sum_{w\in W(C_{d_0/2})}(-1)^{l(w)}X\left(\begin{array}{cc} & \frac{d_0}{2},\ldots,1 \\ w(& \frac{d_0}{2},\ldots,1) \end{array}\right)$.

Let $\mathcal{P} = (d_{2k+1},\ldots,c_0)$ be an orbit in $O(n,\mathbb{C})$, with no two columns are of the same size. Then the character of $X_{\mathcal{P}}$ is of the form

$$X_{\mathcal{P}} \cong (\overbrace{d_{2l+1}}^{+};\overbrace{d_{2l},b_{2l-1}}^{+};\ldots;\overbrace{d_2,d_1}^{+};d_0)$$

where the parts $(\overbrace{d_{2l}, d_{2l-1}}^{+}; \dots ; \overbrace{d_2, d_1}^{+}; d_0)$ *is defined as in the* $Sp(2m,\mathbb{C})$ *case, and* $\overbrace{d_{2l+1}}^{+} = X\left(\begin{array}{c} \frac{d_{2l+1}}{2} - 1, \dots, 0+ \\ \frac{d_{2l+1}}{2} - 1, \dots, 0+ \end{array}\right)$ *if* b_{2d+1} *is even, and* $\overbrace{d_{2l+1}}^{+} = X\left(\begin{array}{c} \frac{d_{2l+1}}{2} - 1, \dots, \frac{1}{2}+ \\ \frac{d_{2l+1}}{2} - 1, \dots, \frac{1}{2}+ \end{array}\right)$ *if* d_{2l+1} *is odd.*

On can easily check the above Proposition holds for Examples 3.6, 3.8 and 3.9.

3.3. **Computing Multiplicities of $X_{\mathcal{Q}}$.** We now compute the multiplicities of the fundamental representations appearing in $R[\mathcal{Q}]$, when no two columns of the orbit Q are of the same size. This is a special case of Theorem 2.3.

Lemma 3.11. *Let* $\overbrace{p,q}^{+}$ *be the character defined as in Proposition 3.10. Then for any fundamental representations* μ_i *in* $G = Sp(2(p+q),\mathbb{C})$ *or* μ_i' *in* $G = SO(2(p+q),\mathbb{C})$*,*

$$[\overbrace{p,q}^{+} : \mu_i] = [Ind_{GL(\frac{p+q}{2})}^{G}(triv) : \mu_i]$$

$$[\overbrace{p,q}^{+} : \mu_i'] = [Ind_{GL(\frac{p+q}{2})}^{G}(triv) : \mu_i']$$

Proof. There are several ways to prove the Lemma, and we present one here. Consider the character formula of $\overbrace{p,q}^{+}$, which is of the form $\overbrace{p,q}^{+} = \sum_i a_i X\left(\begin{array}{c} \phi_i \\ \psi_i \end{array}\right)$. By Frobenius reciprocity

$$[\overbrace{p,q}^{+} : \mu] = \sum_i a_i [X\left(\begin{array}{c} \phi_i \\ \psi_i \end{array}\right) : \mu] = \sum_i a_i [(\phi_i - \psi_i) : \mu|_T]$$

where μ is any fundamental representations of G, and $T \leq G$ is the Cartan subgroup of G. Therefore $\mu|_T$ is just the character of the fundamental representations. Therefore the multiplicities $[\overbrace{p,q}^{+} : \mu_i]$ (or $[\overbrace{p,q}^{+} : \mu_i']$) can be computed, and is equal to.

$$[\overbrace{p,q}^{+} : \mu_i] = \delta_{0i}$$

$$[\overbrace{p,q}^{+} : \mu_i'] = \begin{cases} 1 & \text{if } i \text{ is even} \\ 0 & \text{otherwise} \end{cases}$$

which is precisely the multiplicity of $[Ind_{GL(\frac{p+q}{2})}^{G}(triv) : \mu_i]$ (or $[Ind_{GL(\frac{p+q}{2})}^{G}(triv) : \mu_i']$). Hence the Lemma is proved. □

With the same technique as in the previous proof, we also have the following:

Lemma 3.12. *Let d_0 be an even number, and T be the virtual character*

$$T := \sum_{w \in W(C_{d_0/2})} (-1)^{l(w)} X \left(\begin{array}{c} \frac{d_0}{2}, \dots, 1 \\ w(\quad \frac{d_0}{2}, \dots, 1) \end{array} \right)$$

Then

$$[T : \mu_i] = Ind_{GL(d_0/2,\mathbb{C})}^{Sp(d_0,\mathbb{C})}(triv) = \delta_{i0}$$

in $Sp(d_0, \mathbb{C})$, and

$$[T : \mu_i'] = [Ind_{GL(\frac{d_0}{2},\mathbb{C})}^{SO(d_0,\mathbb{C})}(triv) : \mu_i'] = \begin{cases} 1 & \textit{if } i \textit{ is even} \\ 0 & \textit{otherwise} \end{cases}$$

in $SO(d_0, \mathbb{C})$.

Proposition 3.13. *Let $\mathcal{Q}$ be a nilpotent orbit in $G = Sp(2m, \mathbb{C})$ or $O(n, \mathbb{C})$ such that no two columns are of the same size. Then Theorem 2.3 holds.*

Proof. We first deal with the case of $Sp(2m, \mathbb{C})$. According to the character formula in Proposition 3.10, $X_{\mathcal{Q}}$ has the same virtual character as

$$Ind_{Sp(d_{2l}+d_{2l-1},\mathbb{C})\times\cdots\times Sp(d_2+d_1,\mathbb{C})\times Sp(d_0,\mathbb{C})}^{Sp(2m,\mathbb{C})}(\overbrace{d_{2l}, d_{2l-1}}^{+} \otimes \cdots \otimes \overbrace{d_2, d_1}^{+} \otimes T)$$

Therefore, to compute $[X_{\mathcal{Q}} : \mu_i]$, we need to know:

$$\begin{aligned}
[X_{\mathcal{Q}} : \mu_i] &= [Ind_{Sp(d_{2l}+d_{2l-1},\mathbb{C})\times\cdots\times Sp(d_2+d_1,\mathbb{C})\times Sp(d_0,\mathbb{C})}^{Sp(2m,\mathbb{C})}(\overbrace{d_{2l}, c_{2l-1}}^{+} \otimes \cdots \otimes \overbrace{d_2, d_1}^{+} \otimes T) : \mu_i] \\
&= [\overbrace{d_{2l}, d_{2l-1}}^{+} \otimes \cdots \otimes \overbrace{d_2, d_1}^{+} \otimes T : \wedge^i \mathbb{C}^{2m}|_{Sp(d_{2l}+d_{2l-1},\mathbb{C})\times\cdots\times Sp(d_2+d_1,\mathbb{C})\times Sp(d_0,\mathbb{C})}] \\
&\quad - [\overbrace{d_{2l}, d_{2l-1}}^{+} \otimes \cdots \otimes \overbrace{d_2, d_1}^{+} \otimes T : \wedge^{i-2} \mathbb{C}^{2m}|_{Sp(d_{2l}+d_{2l-1},\mathbb{C})\times\cdots\times Sp(d_2+d_1,\mathbb{C})\times Sp(d_0,\mathbb{C})}] \\
&= [\overbrace{d_{2l}, d_{2l-1}}^{+} \otimes \cdots \otimes \overbrace{d_2, d_1}^{+} \otimes T : \wedge^i \mathbb{C}^{2m}|_{Sp(2z_l,\mathbb{C})\times\cdots\times Sp(2z_1,\mathbb{C})\times Sp(2z_0,\mathbb{C})}] \\
&\quad - [\overbrace{d_{2l}, d_{2l-1}}^{+} \otimes \cdots \otimes \overbrace{d_2, d_1}^{+} \otimes T : \wedge^{i-2} \mathbb{C}^{2m}|_{Sp(2z_l,\mathbb{C})\times\cdots\times Sp(2z_1,\mathbb{C})\times Sp(2z_0,\mathbb{C})}]
\end{aligned}$$

(Recall in Theorem 2.3 that $2z_p = d_{2p} + d_{2p-1}$). Also, the restriction decomposes as $\wedge^i\mathbb{C}^{2m}|_{Sp(2z_l,\mathbb{C})\times\cdots\times Sp(2z_1,\mathbb{C})\times Sp(2z_0,\mathbb{C})} = \bigoplus_{i_0+\cdots+i_l=i}\bigotimes_{p=0}^{l}\wedge^{i_p}\mathbb{C}^{2z_p}$. So

$$\begin{aligned}
[X_{\mathcal{Q}} : \mu_i] &= [\overbrace{d_{2l}, d_{2l-1}}^{+} \otimes \cdots \otimes \overbrace{d_2, d_1}^{+} \otimes T : \bigoplus_{i_0+\cdots+i_l=i}\bigotimes_{p=0}^{l}\wedge^{i_p}\mathbb{C}^{2z_p}] \\
&\quad - [\overbrace{d_{2l}, d_{2l-1}}^{+} \otimes \cdots \otimes \overbrace{d_2, d_1}^{+} \otimes T : \bigoplus_{i_0+\cdots+i_l=i-2}\bigotimes_{p=0}^{l}\wedge^{i_p}\mathbb{C}^{2z_p}] \\
&= \sum_{i_0\cdots+i_l=i}[T : \wedge^{i_0}\mathbb{C}^{2z_0}]\prod_{p=1}^{l}[\overbrace{d_{2p}, d_{2p-1}}^{+} : \wedge^{i_p}\mathbb{C}^{2z_p}] \\
&\quad - \sum_{i_1\cdots+i_l=i-2}[T : \wedge^{i_0}\mathbb{C}^{2z_0}]\prod_{p=1}^{l}[\overbrace{d_{2p}, d_{2p-1}}^{+} : \wedge^{i_p}\mathbb{C}^{2z_p}]
\end{aligned}$$

On the other hand, by Lemma 3.11 and Lemma 3.12,

$$\begin{aligned}
[\overbrace{d_{2p}, d_{2p-1}}^{+} : \wedge^{i}\mathbb{C}^{2z_p}] &= [\overbrace{d_{2p}, d_{2p-1}}^{+} : \mu_i \oplus \mu_{i-2} \oplus \mu_{i-4} \oplus \dots] \\
&= \sum_{0\leq i-2q\leq i}[\overbrace{d_{2p}, d_{2p-1}}^{+} : \mu_{i-2q}] \\
&= \sum_{0\leq i-2q\leq i}[Ind_{GL(z_p,\mathbb{C})}^{Sp(2z_p,\mathbb{C})}(triv) : \mu_{i-2q}] \\
&= [Ind_{GL(z_p,\mathbb{C})}^{Sp(2z_p,\mathbb{C})}(triv) : \wedge^{i}\mathbb{C}^{2z_p}]
\end{aligned}$$

and

$$[T : \wedge^{i}\mathbb{C}^{2z_0}] = [Ind_{GL(z_0,\mathbb{C})}^{Sp(2z_0,\mathbb{C})}(triv) : \wedge^{i}\mathbb{C}^{2z_0}]$$

Consequently,

$$\begin{aligned}
[X_{\mathcal{Q}} : \mu_i] &= \sum_{i_0\cdots+i_l=i}\prod_{p=0}^{l}[Ind_{GL(z_p,\mathbb{C})}^{Sp(2z_p,\mathbb{C})}(triv) : \wedge^{i_p}\mathbb{C}^{2z_p}] \\
&\quad - \sum_{i_0\cdots+i_l=i-2}\prod_{p=0}^{l}[Ind_{GL(z_p,\mathbb{C})}^{Sp(2z_p,\mathbb{C})}(triv) : \wedge^{i_p}\mathbb{C}^{2z_p}]
\end{aligned}$$

Reversing the process, we get

$$[X_{\mathcal{Q}} : \mu_i] = [Ind_{GL(z_l,\mathbb{C})\times\cdots\times GL(z_0,\mathbb{C})}^{Sp(2m,\mathbb{C})}(triv \otimes \cdots \otimes \ triv) : \mu_i]$$

Then the result follows by Frobenius reciprocity and induction on k.
For the $O(n,\mathbb{C})$ case, $X_{\mathcal{P}}$ has the same virtual character as

$$Ind_{O(d_{2l+1},\mathbb{C})\times SO(d_{2l}+d_{2l-1},\mathbb{C})\times\cdots\times SO(d_2+d_1,\mathbb{C})\times SO(d_0,\mathbb{C})}^{O(n,\mathbb{C})}(triv \otimes \overbrace{d_{2l}, d_{2l-1}}^{+} \otimes \cdots \otimes \overbrace{d_2, d_1}^{+} \otimes T)$$

which, by the same argument as before, gives

$$[X_{\mathcal{P}} : \mu_i'] = [Ind_{O(d_{2l+1},\mathbb{C})\times GL(z_l,\mathbb{C})\times\cdots\times GL(z_0,\mathbb{C})}^{O(n,\mathbb{C})}(triv \otimes \cdots \otimes triv) : \mu_i']$$

the result follows again by Frobenius reciprocity and induction on k. □

3.4. **Proof of Theorem 2.3.** With Proposition 3.13 on hand, it is now easy to see how Theorem 2.3 works. Recall that

$$X_{\mathcal{O}} := Ind_{G(\mathcal{Q},\mathbb{C})\times GL(Y)}^{G(2m,\mathbb{C})}(X_{\mathcal{Q}} \otimes triv \otimes \cdots \otimes triv)$$

By Proposition 3.12 and induction in stages, for any fundamental representations μ,

$$[X_{\mathcal{O}} : \mu] = [Ind_{GL(Y)\times GL(Z,\mathbb{C})}^{Sp(2m,\mathbb{C})}(triv \otimes \cdots \otimes triv) : \mu]$$

(Recall the notations in Theorem 2.3). Similarly, if $\mathcal{P}$ is any orbit in $O(n,\mathbb{C})$,

$$[X_{\mathcal{P}} : \mu] = [Ind_{GL(Y)\times O(d_{2l+1},\mathbb{C})\times GL(Z,\mathbb{C})}^{O(n,\mathbb{C})}(triv \otimes \cdots \otimes triv) : \mu]$$

the Theorem follows in exactly the same fashion as in Proposition 3.12. □

Remark 3.14. *In fact, the model given by Barbasch in* [4] *does not only deal with nilpotent orbits, but also some finite covers of nilpotent orbits* $\widetilde{\mathcal{O}} \twoheadrightarrow \mathcal{O}$. *One can also compute the multiplicities of finite dimensional irreducible representations in* $R[\widetilde{\mathcal{O}}]$ *using the same technique.*

4. Proof of Corollary 2.5

With the algoritms above, we can now state and prove another criterion of the normality of $\overline{\mathcal{O}}$ by computing the multiplicities of irreducible finite-dimensional representations in $X_{\mathcal{O}} \cong R[\mathcal{O}]$ (Theorem 3.2). Note that by the hypothesis in Theorem 2.5, the longest column of the orbit in $O(n,\mathbb{C})$ is not involved. So we can just focus ourselves on the case of $Sp(2m,\mathbb{C})$.

Lemma 4.1. *Let* $\mathcal{O} = (c_{2k}, c_{2k-1}, \ldots, c_0)$ *be a nilpotent orbit in* $G = Sp(2m,\mathbb{C})$, *and* μ *be* ***any*** *finite dimensional irreducible representation of* G, *then*

$$[R[\overline{\mathcal{O}}] : \mu] \leq [R[\mathcal{O}^{\sharp}] : \mu]$$

where $\mathcal{O}^{\sharp} = (\frac{c_{2k}+c_{2k-1}}{2}, \frac{c_{2k}+c_{2k-1}}{2}, \frac{c_{2k-2}+c_{2k-3}}{2}, \frac{c_{2k-2}+c_{2k-3}}{2}, \ldots, \frac{c_2+c_1}{2}, \frac{c_2+c_1}{2}, c_0)$

Proof. Note that by the Kraft-Procesi criterion (Theorem 1.6), $\overline{\mathcal{O}^\sharp}$ is normal. Therefore Corollary 2.2 says $R[\mathcal{O}^\sharp] = R[\overline{\mathcal{O}^\sharp}]$. On the other hand, note that $\overline{\mathcal{O}^\sharp} \supset \overline{\mathcal{O}}$. Consequently, we have a G-module surjection

$$R[\mathcal{O}^\sharp] = R[\overline{\mathcal{O}^\sharp}] \twoheadrightarrow R[\overline{\mathcal{O}}]$$

and hence $[R[\overline{\mathcal{O}}] : \mu] \leq [R[\overline{\mathcal{O}^\sharp}] : \mu]$ for any finite dimensional G-representations μ. However, the latter term is equal to $[R[\mathcal{O}^\sharp] : \mu]$. Hence the result follows. □

From results in the last Section (or from [10] directly), one can find out the multiplicities of $X_{\mathcal{O}^\sharp}$. And now we are in the position to prove Corollary 2.5:

4.1. **Proof of Corollary 2.5.** One direction is easy - if $\overline{\mathcal{O}}$ is normal, then $R[\overline{\mathcal{O}}] = R[\mathcal{O}]$ as G-modules, hence $[R[\overline{\mathcal{O}}] : \mu_i] = [R[\mathcal{O}] : \mu_i]$ for all i.
Now suppose $\overline{\mathcal{O}}$ is not normal, and let $\mathcal{O}^\sharp$ as in last Section. Then we obtain a new set of integers $\{x_i\}$ computing the multiplicities of $X_{\mathcal{O}^\sharp}$. By the Kraft-Procesi criterion (Theorem 1.6), the two sets of integers $\{x_i\}$ (from $\mathcal{O}^\sharp$) and $\{w_i\}$ (from $\mathcal{O}$) are different. More precisely, for $m := \min\{i | x_i \neq w_i\}$, $x_m < w_m$. By the algorithm in Theorem 2.3, this implies the existence of an $i > 0$ such that $[R[\mathcal{O}^\sharp] : \mu_i] < [R[\mathcal{O}] : \mu_i]$. Now Lemma 4.1 says $[R[\overline{\mathcal{O}}] : \mu_i] \leq [R[\mathcal{O}^\sharp] : \mu_i]$ for all i, and consequently the theorem follows. □

Example 4.2. *Going back to Example 2.4. Let* $\mathcal{O} = (8,6,6,4,4,2,2)$ *in* $Sp(32,\mathbb{C})$*. Following Theorem 1.6 (Kraft-Procesi criterion), its closure is not normal.*
By Theorem 2.3, the w_i*'s are* $\{2,4,4,6\}$*. Now* $\mathcal{O}^\sharp = (7,7,5,5,3,3,2)$*, and the* x_i*'s are* $\{1,3,5,7\}$*. We therefore have the multiplicities as follows:*

i	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
$[R[\mathcal{O}^\sharp] : \mu_i]$	1	0	3	0	5	0	7	0	8	0	8	0	7	0	5	0	2
$[R[\mathcal{O}] : \mu_i]$	1	0	3	0	6	0	9	0	12	0	13	0	12	0	8	0	3

The discrepancies of the two rows of numbers reflects the non-normality of $\overline{\mathcal{O}}$*.*

Let $\mathcal{P} = (7,5,3,3,1)$ *in* $O(19,\mathbb{C})$*. Then* $\mathcal{P}^\sharp = (7,4,4,2,2)$*, the* w_i*'s are* $\{6,6\}$*, and the* x_i*'s are* $\{4,8\}$*. We therefore have the multiplicities of* μ'_{2i} *as follows:*

i	0	2	4	6	8	10	12	14	16	18
$[R[\mathcal{P}^\sharp] : \mu'_i]$	1	2	3	3	3	2	1	0	0	0
$[R[\mathcal{P}] : \mu'_i]$	1	2	3	4	3	2	1	0	0	0

5. Final Remarks

By Lemma 4.1, there is an upper bound of the multiplicities of $R[\overline{\mathcal{O}}]$ by that of $R[\mathcal{O}^\sharp]$. In fact, we have the following

Conjecture 5.1. *Let $\mathcal{O}$ be a classical nilpotent orbit in G, and μ is any irreducible, finite dimensional representation of G. Then the multiplicities $[R[\overline{\mathcal{O}}] : \mu]$ can be computed. In particular, if μ is a fundamental representation, then*

$$[R[\overline{\mathcal{O}}] : \mu] = R[\mathcal{O}^\sharp] : \mu]$$

The author knows of no counter-example to the above conjecture.

References

[1] Anker, J-P. and Orsted, B., *Lie theory: Lie algebras and representations*, Birkhauser, 2004

[2] Barbasch, D. and Vogan, D., *Unipotent Representations of Complex Semisimple Groups*, Annals of Mathematics **121**, No.1, 41-110, 1985

[3] Barbasch, D., *The Unitary Dual for Complex Classical Lie Groups*, Invent. Math. **96**, 103-176, 1989

[4] Barbasch, D., *Regular Functions on Covers of Nilpotent Coadjoint Orbits* `http://arxiv.org/abs/0810.0688v1`, 2008

[5] Brylinski, R., *Dixmier Algebras for Classical Complex Nilpotent Orbits via Kraft-Procesi Models I*, The orbit method in geometry and physics: in honor of A.A. Kirillov, Birkhauser, 2003

[6] Collingwood, D. and McGovern, W., *Nilpotent orbits in semisimple Lie algebras*, Van Norstrand Reinhold Mathematics Series, 1993

[7] Eisenbud, D. *Commutative Algebra: with a View Toward Algebraic Geometry*, Graduate Texts in Mathematics, Springer-Verlag, 1995

[8] Humphreys, J. *ntroduction to Lie Algebras and Representation Theory*, Graduate Texts in Mathematics, Springer-Verlag, 1973

[9] Kraft, H. and Procesi, C., *On the Geomery of Conjugacy Classes in Classical Groups*, Comment. Math. Helvetici **57**, 539-602, 1982

[10] McGovern, W., *Rings of regular functions on nilpotent orbits and their covers*, Invent. Math. **97**, 209-217, 1989

[11] McGovern, W., *Completely Prime Maximal Ideals and Quantization*, Memoirs of the American Mathematical Society **519**, 1994

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA NY 14853, USA
E-mail: `kywong@math.cornell.edu`
Website: `http://www.math.cornell.edu/`∼ `kywong`